

April 1, 2014

## Snowpack Remains Way Above Normal

The USDA Natural Resources Conservation Service (NRCS) Kremmling Field Office snow surveyors Mark Volt, Noah Bates took the April 1 snow survey measurements during the last days of March. Snowpack for Middle Park and the upper Colorado River Basin stands at 144% of average. We were at 79% last April 1<sup>st</sup>. and 135% in the high year of 2011 and 62% in the drought year of 2002.

Snowpack in the mountains above Middle Park now ranges from 114% to 206% of the 30-year average. The Granby snow course near C Lazy U Ranch set a new record high for the second month in a row. It has been read since 1949. Snow density is averaging 31%, which means that for a foot of snow there are 3.72 inches of water. Irrigators, towns, river runners and other water users can expect higher than normal river levels this summer. Reservoir storage remains higher than last year. From this point on, spring runoff will be highly dependent on melting conditions (i.e., temperature and wind), as well as spring snow accumulation and/or rainfall.

Reported average readings for the major river basins in Colorado are as follows: Colorado River Basin 1287%; Gunnison River Basin, 111%; South Platte River Basin, 135%; Yampa and White River Basins, 124%; Arkansas River Basin, 101%; Upper Rio Grande Basin, 84%; San Miguel, Dolores, Animas, and San Juan River Basins 90%; and Laramie and North Platte River Basins, 139%.

Most of the snow courses around Middle Park have been read since the 1940s. Snow course readings are taken at the end of each month, beginning in January and continuing through April. March is historically the snowiest month, and the April 1 readings are the most critical for predicting runoff and summer water supplies, as most of our high country snowpack peaks around that time. Manual snow courses will be read for the final time this year at the end of April.

For further information, including real-time snow and precipitation data for SNOTEL (automated Snow Telemetry) sites, visit <http://www.co.nrcs.usda.gov/snow/index.html>.



# NEWS RELEASE

NRCS Kremmling Field Office snow survey for April 1, 2014, compared to long-term average.								
Snow course or SNOTEL	Last year		This year		30-year average (1981 - 2010)		Percent of average	
	Snow depth	Moisture content	Snow depth	Moisture content	Snow depth	Moisture content	Snow depth	Moisture content
	------(inches)-----						-----(%)------	
Arapaho Ridge st		17.6		26.1		19.8		132%
Berthoud Summit st		15.3		22.7		17.9		127%
Buffalo Park st		12.3		20.4		10.4		196%
Columbine st		19.6		34.6		22.1		157%
Copper Mountain st		10.1		19.8		13.2		150%
Corral Creek sc	35	9.2		not read	45	14.6	77	#VALUE!
Elliot Ridge st		15.4						
Fremont Pass st		12.6		19		14.8		128%
Gore Pass sc	35	10.2	45	15.3	34	10.6	132	144%
Granby sc	22	5.4	45	14.8	26	7.2	205	206%
Grizzly Peak st		12.8		23.6		15.8		149%
Jones Pass st		11.7		18		12.9		140%
Lake Irene st		16.3		31.2		23.5		133%
Lynx Pass st		9.8		13.4		10.8		124%
Middle Fork Camp sc	37	9.9	42	13.7	35	10.1	136	136%
Phantom Valley st		7.2		12.1		9.1		133%
Stillwater Creek st		5.2		not read		7		#VALUE!
Summit Ranch st		7.5		13.8		10.8		128%
Willow Creek Pass sc	37	9.7	49	14.3	44	12.5	111	114%
<i>Average</i>								<b>144%</b>

# NEWS RELEASE

## NRCS Kremmling Field Office snow survey April 1 moisture content records.

Snow course or SNOTEL (SC or ST)	Highest Apr. 1 moisture content			Lowest Apr. 1 moisture content		
	(inches)	(%)	(year)	(inches)	(%)	(year)
Arapaho Ridge ST (read since 2003)	26.9	117%	2008	14.8	65%	2010
Berthoud Summit ST	25.3	138%	1996	10.9	59%	2002
Buffalo Park ST (read since 1996)	17.7	123%	1996	8.0	56%	2002
Columbine ST	37.9	152%	1984	12.0	48%	1981
Copper Mountain ST	18.1	129%	2008	7.9	56%	1981
Corral Creek SC (read since 1995)	16.9	116%	1996	8.0	55%	2002
Fremont Pass ST	21.9	135%	1978	10.0	62%	1966
<b>Gore Pass SC</b>	<b>16.6</b>	<b>157</b>	<b>2011</b>	4.3	41%	1966
Granby SC	14.8	206%	2014	0.0	0%	2004
Grizzly Peak ST	27.7	151%	1996	8.7	47%	1981
Jones Pass ST (read since 2000)	19.2	119%	2006	9.7	60%	2002
Lake Irene ST	37.6	146%	1962	11.8	46%	1977
Lynx Pass ST	20.8	163%	1962	6.0	47%	1977
Middle Fork Campground SC	17.0	168%	1996	5.8	57%	1981
Phantom Valley ST	14.9	160%	1996	1.1	12%	2004
Stillwater Creek ST (read since 1986)	12.8	164%	1965	1.1	14%	2012
Summit Ranch ST	17.0	153%	1996	5.9	53%	2012
Willow Creek Pass SC	20.5	164%	1952	6.6	53%	2012