

NEWS RELEASE

U.S. Department of Agriculture • Natural Resources Conservation Service • 106 S 2nd Street, PO Box 265 Kremmling, CO 80459-0265 • Phone: (970)724-3456 • Fax: (970)724-0807 • http://www.co.nrcs.usda.gov

March 1, 2014

Snowpack in Middle Park Remains Way Above Normal

The USDA Natural Resources Conservation Service (NRCS) Kremmling Field Office snow surveyors Mark Volt and Noah Bates took the March 1st. snow survey measurements during the last days of February.....And even with the recent warm weather, the survey shows that we are in good shape!

Snowpack in the high elevation mountains above Middle Park now ranges from 115% to 194% of the 30-year average, with the overall average for Middle Park at 141%. New record highs were set at both the Granby and Buffalo Park Snow courses. Last year at this time we were only at 75% of average. Snow density is averaging 28%, which means that for one foot of snow there is only 3.3 inches of water, which is pretty normal for March 1st.

All Colorado River basins are reporting above average snowpack except for southwestern Colorado which is 5%-15% below average. Reported readings for the major river basins in Colorado are as follows: Colorado River Basin 137%; Gunnison River Basin, 114%; South Platte River Basin,145%; Yampa and White River Basins, 122%; Arkansas River Basin, 103%; Upper Rio Grande Basin, 85%; San Miguel, Dolores, Animas, and San Juan River Basins 95%; and the Laramie and North Platte River Basins, 132% of average for this time of year.

Most of the snow courses around Middle Park have been read since the 1940s. Snow course readings are taken at the end of each month, beginning in January and continuing through April. March is historically the snowiest month, and the April 1 readings are the most critical for predicting runoff and summer water supplies, as most of our high country snowpack peaks around that time.

For further information, including real-time snow and precipitation data for SNOTEL (automated Snow Telemetry) sites, visit http://www.co.nrcs.usda.gov/snow/index.html.



NEWS RELEASE

NRCS Kremmling Field Office snow survey for March 1, 2014, compared to long-term average.

S. SNOTEL	Last year		This year		30-year average (1971 - 2000)		Percent of average	
Snow course or SNOTEL (SC or ST)	Snow depth	Moisture content	Snow depth	Moisture content	Snow depth	Moisture content	Snow depth	Moisture content
			(i	(%)				
Arapaho Ridge ST		15.0		21.6		16.3		133%
Berthoud Summit ST		11.4		19.5		13.5		144%
Buffalo Park ST		12.0		17.1		10.8		188%
Columbine ST		17.7				19.4		147%
Copper Mountain ST		9.2		16.8		10.6		158%
Corral Creek SC	36	10.7	54	15.8	48	13.1	113%	121%
Elliot Ridge ST	11.6			16.6	New site		New site	
Fremont Pass ST		10.2		15.3		11.6		132%
Gore Pass SC	43	9.6	46	12.6	34	9.2	135%	137%
Granby SC	40	6.8	42	12.4	27	6.4	156%	194%
Grizzly Peak ST		11.0		18.8		12.8		147%
Jones Pass ST		8.3		13.8		9.5		145%
Lake Irene ST		15.2		24.4		18.9		130%
Lynx Pass ST		7.7		11.3		8.7		130%
Middle Fork Camp SC	27	6.9	39	11.2	35	8.7	112%	129%
Phantom Valley ST		6.7		10.2		8.1		126%
Stillwater Creek ST		5.6		9.9		6.5		152%
Summit Ranch ST		7.4		10		8.5		118%
Willow Creek Pass SC	34	7.3	43	10.7	40	9.3	108%	115%
Average for Middle Park								141%





United States Department of Agriculture Natural Resources Conservation Service NEWS RELEASE

NRCS Kremmling Field Office snow survey March 1 moisture content records.

Snow course or SNOTEL (SC or ST)	Highest March. 1 moisture content			Lowest March. 1 moisture content			
	(inches)	(%)	(year)	(inches)	(%)	(year)	
Arapaho Ridge ST (read since 2003)	23.5	125%	2006	12.8	68%	2010	
Berthoud Summit ST	20.3	142%	1996	7.5	52%	1981	
Buffalo Park ST (read since 1996)	17.1	188%	2014	6.6	61%	2002	
Columbine ST	40.1	183%	1984	8.2	37%	1981	
Copper Mountain ST	18.9	169%	1996	4.8	43%	1981	
Corral Creek SC (read since 1995)	18.0	137%	1997	6.6	50%	2010	
Fremont Pass ST	18.8	146%	2006	6.4	50%	1981	
Gore Pass SC	14.3	155%	1962	2.6	28%	1981	
Granby SC (New Record High)	12.4	194%	2014	1.8	28%	1981	
Grizzly Peak ST	23.7	173%	1996	6.1	45%	1981	
Jones Pass ST (read since 2000)	18.1	141%	2006	8.5	66%	2010	
Lake Irene ST	32.3	154%	1956	7.1	34%	1977	
Lynx Pass ST	17.5	172%	1962	2.8	27%	1981	
Middle Fork Campground SC	13.9	160%	1996	3.0	34%	1981	
Phantom Valley ST	12.0	143%	1996	3.4	40%	1977	
Stillwater Creek ST (read since 1986)	11.4	158%	1997	2.9	40%	1981	
Summit Ranch ST	15.4	181%	1996	5.0	59%	1981	
Willow Creek Pass SC	20.4	192%	1952	3.8	36%	1977	